arXiv:1303.5326v1 [quant-ph] 21 Mar 2013

Greenberger-Horne-Zeilinger theorem for N-partite qu$D$its

Junghee Ryu,[1, 2] Changhyoup Lee,[3, 2] Marek Żukowski,[1] and Jinhyoung Lee[2, 4]

[1] *Institute of Theoretical Physics and Astrophysics, University of Gdańsk, 80-952 Gdańsk, Poland*
[2] *Department of Physics, Hanyang University, Seoul 133-791, Korea*
[3] *Centre for Quantum Technologies, National University of Singapore, 3 Science Drive 2, Singapore 117543*
[4] *Center for Macroscopic Quantum Control, Seoul National University, Seoul, 151-747, Korea*

(Received March 22, 2013)

We generalize Greenberger-Horne-Zeilinger (GHZ) theorem to an arbitrary number of D-dimensional systems. Contrary to conventional approaches, we use concurrent observables, which are incompatible but still have a common eigenstate. We begin with the theorem for 4 systems of a dimension divisible by 3, and discuss its extension to N systems of an arbitrary dimension. The GHZ theorem can be proved as long as N is not divisible by all non-unit divisors of D, smaller than N.

I. INTRODUCTION

Bell's theorem says there exists quantum correlations between separated observers which cannot be simulated by local realistic theories [1–3]. Using a statistical inequality, Bell showed that any local realistic theories are incompatible with quantitative predictions of quantum mechanics for a singlet state of two spin-1/2 particles [1]. Even though Bell's theorem was studied mostly in terms of statistical inequalities, different version of it *without inequalities*, was also shown for a multiqubit systems by Greenberger, Horne, and Zeilinger (GHZ) [4]. They derived an *all-versus-nothing* contradiction based on perfect correlations of the so-called GHZ states. This leads to a direct refutation of EPR ideas on relation between locality and elements of reality with quantum mechanics.

Extending Bell's theorem is important not only for a deeper understanding of foundations of quantum mechanics. It allows developing new applications in quantum information processing, such as quantum cryptography, secret sharing, quantum teleportation, reduction of communication complexity, quantum key distribution and random numbers generation [5–9]. A Bell-type test is also regarded as a fundamental method to verify entanglement among subsystems. Therefore, search for extensions of Bell's theorem to more complex systems such as multipartite and/or high-dimensional systems is an important task in quantum information science [3].

An all-versus-nothing test, that we call GHZ theorem, has been generalized to higher dimensional systems. For the sake of convenience, we shall use the tuple (N, M, D) to denote N parties, M measurements for each party and D distinct outcomes for each measurement. In Ref. [10], GHZ theorem was derived for a $(D+1, 2, D)$ problem. A probabilistic but conclusive GHZ-like test was shown for $(D, 2, D)$ in Ref. [11]. The (N, M, D) problem for odd $N > D$, $M = 2$, and even D was studied using operator relations by Cerf *et al.* [12]. Recently, Lee *et al.* employed incompatible composite observables for arbitrary $(\text{odd } N, 2, \text{even } D)$ problems [13]. The conventional approaches are based on compatible observables.

We generalize the GHZ theorem to 3 or more D-dimensional systems. To this end, we employ concurrent composite observables. Such observables are mutually incompatible but still have a common eigenstate, here a generalized GHZ state. They can be realized by using multiport beam splitters and phase shifters, as it is shown in Refs. [10, 13, 14]. We illustrate our principal idea with four $3d$-dimensional systems, and provide a systematic method to extend it to three or more D-dimensional systems. As non-unit divisors of D play a crucial role, one can show a GHZ-type contradiction as long as N is not divided by all non-unit divisors of D, which are smaller than N. We also show that our approach can reproduce the previous results [4, 10, 12, 13] and it can lead to a general GHZ theorem for N qu$D$its.

II. CONCURRENT OBSERVABLES AND INVARIANCE CONDITION

Some sets of observables have a common eigenstate. If a system is prepared in the eigenstate, the measurement results for such observables are concurrently appearing with certainty. Such observables are called "concurrent" [13]. For a quantum system of dimension D, greater than two, consider two Hermitian operators $\hat{A}$ and $\hat{B}$, such that $\hat{A} = a|\psi\rangle\langle\psi| + \hat{A}'$ and $\hat{B} = b|\psi\rangle\langle\psi| + \hat{B}'$. The state $|\psi\rangle$ is then a common eigenstate of both observables as $\hat{A}(\hat{B})|\psi\rangle = a(b)|\psi\rangle$, even if $[\hat{A}, \hat{B}] = [\hat{A}', \hat{B}'] \neq 0$. Note that compatible observables are clearly concurrent.

To construct concurrent observables, in a Hilbert space $\mathcal{H}$, one can use the method of Ref. [13]. Consider a unitary operator $\hat{U}$, which is of the form of $\hat{U} = e^{i\phi}|\psi\rangle\langle\psi| + \hat{U}^{\perp}$ with $\hat{U}^{\perp}|\psi\rangle = 0$. Here $\hat{U}^{\perp}$ is a unitary operator on a space $\mathcal{H}^{\perp}$ which is defined by the requirement $\mathcal{H} = \mathcal{H}^{\psi} \oplus \mathcal{H}^{\perp}$, where $\mathcal{H}^{\psi}$ is the one-dimensional space containing $|\psi\rangle$. Every such unitary operator leaves the state $|\psi\rangle$ unchanged, up to a global phase. If the state $|\psi\rangle$ satisfies $\hat{A}|\psi\rangle = \lambda|\psi\rangle$, then all transformed operators $\hat{B}_U = \hat{U}\hat{A}\hat{U}^{\dagger}$ are concurrent with $\hat{A}$.

We shall use the following local observables. Consider a measurement in D-dimensional Hilbert space represented by $\hat{A} = \sum_{n=0}^{D-1} \omega^n |n\rangle_{\mathrm{AA}}\langle n|$, where $\omega =$

$\exp(2\pi i/D)$. The operator $\hat{A}$ is unitary, however one can uniquely relate it with a Hermitian observable $\hat{H}$, by requiring $\hat{A} = \exp(i\hat{H})$. Therefore the complex eigenvalues of $\hat{A}$, which could be called a unitary observable, can be associated with the measurement results, denoted by real eigenvalues of $\hat{H}$ [12, 13, 15]. Such a unitary representation leads to a simplification of the mathematics without changing any physical results.

Consider N qu$D$its in a generalized GHZ state,

$$|\psi\rangle = \frac{1}{\sqrt{D}} \sum_{n=0}^{D-1} \underbrace{|n\rangle \otimes \cdots \otimes |n\rangle}_{N}, \tag{1}$$

where $|n\rangle$ denotes a basis state for a qu$D$it. Let us consider a unitary operation in the form of $\hat{U} = \bigotimes_{k=1}^{N} \hat{P}_k(f_k)$ with phase shifter $\hat{P}_k = \sum_{n=0}^{D-1} \omega^{f_k(n)} |n\rangle\langle n|$. If "phases" $f_k(n)$ satisfy a condition

$$\sum_{k=1}^{N} f_k(n) \equiv 0 \mod D, \tag{2}$$

for each n, then the unitary operator $\hat{U}$ leaves the GHZ state $|\psi\rangle$ invariant.

Consider now a composite observable $\hat{X}^{\otimes N} \equiv \hat{X} \otimes \hat{X} \otimes \cdots \otimes \hat{X}$, where the local observable $\hat{X}$ is defined by applying quantum Fourier transformation $\hat{F}$ on a reference unitary observable $\hat{Z} = \sum_{n=0}^{D-1} \omega^n |n\rangle\langle n|$, that is $\hat{X} = \hat{F}\hat{Z}\hat{F}^\dagger$. An eigenvector of $\hat{X}$ associated with the eigenvalue ω^n is given by

$$|n\rangle_{\mathrm{x}} = \hat{F}|n\rangle = \frac{1}{\sqrt{D}} \sum_{m=0}^{D-1} \omega^{nm} |m\rangle. \tag{3}$$

With the standard basis set $\{|n\rangle\}$, the observable $\hat{X}$ can put as by $\hat{X} = \sum_{n=0}^{D-1} |n\rangle\langle n+1|$, where $|n\rangle \equiv |n \mod D\rangle$. The generalized GHZ state $|\psi\rangle$ is an eigenstate of the composite observable $\hat{X}^{\otimes N}$ with the unity eigenvalue: $\hat{X}^{\otimes N}|\psi\rangle = \hat{X} \otimes \hat{X} \otimes \cdots \otimes \hat{X}|\psi\rangle = |\psi\rangle$.

We can construct a set of concurrent composite observables by applying the unitary operation $\hat{U} = \bigotimes_{k=1}^{N} \hat{P}_k$ to $\hat{X}^{\otimes N}$. Consider the phases $f_k(n) = \alpha_k n$ with rational numbers α_k. If the phases $f_k(n)$ satisfy the invariance condition (2), the transformed composite observable $\hat{U}\hat{X}^{\otimes N}\hat{U}^\dagger = \hat{X}(\alpha_1) \otimes \hat{X}(\alpha_2) \otimes \cdots \otimes \hat{X}(\alpha_N)$ is concurrent with $\hat{X}^{\otimes N}$, i.e., $\hat{U}\hat{X}^{\otimes N}\hat{U}^\dagger|\psi\rangle = |\psi\rangle$.

For each eigenvalue ω^n, the eigenvector of the local observable $\hat{X}(\alpha)$ is given by applying the phase shifter $\hat{P}$ on $|n\rangle_{\mathrm{x}}$ as

$$|n\rangle_\alpha = \hat{P}|n\rangle_{\mathrm{x}} = \frac{1}{\sqrt{D}} \sum_{m=0}^{D-1} \omega^{(n+\alpha)m} |m\rangle. \tag{4}$$

The observable $\hat{X}(\alpha)$ can be written in the standard basis set $\{|n\rangle\}$ as

$$\hat{X}(\alpha) = \omega^{-\alpha} \left(\sum_{n=0}^{D-2} |n\rangle\langle n+1| + \omega^{\alpha D} |D-1\rangle\langle 0| \right). \tag{5}$$



FIG. 1. The invariant condition for 4-partite system. Each party applies a local phase shifter $\hat{P}_k$ with phase $f_k(n)$ to the generalized GHZ state $|\psi\rangle$. If the sum of the local phases $f_k(n)$ is congruent to zero modulo D, i.e., $f_1(n) + f_2(n) + f_3(n) + f_4(n) \equiv 0 \mod D$, then the state $|\psi\rangle$ transformed by the phase shifters is unchanged.

Note that if α is an integer, the measurement basis set $\{|n\rangle_\alpha\}$ of $\hat{X}(\alpha)$ will be the same as $\{|n\rangle_{\mathrm{x}}\}$ of $\hat{X}$ except the ordering, i.e. $|n\rangle_\alpha = |n+\alpha\rangle_{\mathrm{x}}$, and then $\hat{X}(\alpha) = \omega^{-\alpha}\hat{X}$. That is, the observable $\hat{X}(\alpha)$ is equivalent to $\hat{X}$, up to a phase factor $\omega^{-\alpha}$. Let $|n\rangle_\alpha$ be the eigenstate of $\hat{X}(\alpha)$ associated with eigenvalue ω^n, and $|m\rangle_\beta$ the eigenstate of $\hat{X}(\beta)$ associated with ω^m. If and only if α differs from β by an integer, then two measurement bases satisfy $|{}_\alpha\langle n|m\rangle_\beta|^2 = \delta_D(\gamma)$, where $\gamma = m - n + \beta - \alpha$. Here $\delta_D(\gamma) = 1$ if γ is congruent to zero modulo D and otherwise $\delta_D(\gamma) = 0$. That is, unless $\beta - \alpha$ is an integer, two local observables $\hat{X}(\alpha)$ and $\hat{X}(\beta)$ are inequivalent.

III. GENERALIZED GHZ THEOREM

A. Four-partite systems

We illustrate our idea by considering first a four qu$D$it system. Already this case goes significantly beyond the previous studies [10, 12, 13].

Take a 4 qu$D$it GHZ state (D is an integral multiple of 3)

$$|\psi\rangle = \frac{1}{\sqrt{D}} \sum_{n=0}^{D-1} |n, n, n, n\rangle. \tag{6}$$

The qu$D$its are distributed to four sufficiently separated parties. Each party performs one of two non-degenerate local measurements on his or her qu$D$it, each of which produces distinguishable D outcomes. The eigenvalues of the observables are of the form ω^k.

Consider such a measurement, $\hat{M}_i$ for the i-th party. One can denote the probability that the parties obtain results ω^{m_i}, where m_i is an integer, by $\mathcal{P}(m_1, m_2, m_3, m_4)$, and define a correlation function $E_{\mathrm{QM}}(M_1, M_2, M_3, M_4) = \langle\psi|\hat{M}_1 \otimes \hat{M}_2 \otimes \hat{M}_3 \otimes \hat{M}_4|\psi\rangle$, the quantum average of products of the measurement results:

$$\sum_{m_1=0}^{D-1}\cdots\sum_{m_4=0}^{D-1}\omega^{m_1+m_2+m_3+m_4}\mathcal{P}(m_1,...,m_4). \tag{7}$$

If all measurements are $\hat{X}$, that is each $\hat{M}_i=\hat{X}$, since

$$\hat{X}\otimes\hat{X}\otimes\hat{X}\otimes\hat{X}|\psi\rangle=|\psi\rangle, \tag{8}$$

one has $E_{\mathrm{QM}}(X,X,X,X)=1$. We have a perfect GHZ correlation. If arbitrary three parties know their own outcomes ω^{x_i} of measurements $\hat{X}$, then they can predict with certainty the remaining party's outcome: We will denote such a perfect correlation by

$$C_{\mathrm{QM}}(x_1+x_2+x_3+x_4\equiv 0). \tag{9}$$

Of course, the sum is taken modulo D; such a convention is used below in all formulas.

We can now construct concurrent composite observables from the observable $\hat{v}_0=\hat{X}^{\otimes 4}$, by applying a unitary operator $\hat{U}_1=\hat{P}_1\otimes\hat{P}_2^{\otimes 3}$, with the phase shifters $\hat{P}_k$ of phases $f_1(n)=(D-1)n$ and $f_2(n)=n/3$. One of the new observables is $\hat{v}_1\equiv\hat{U}_1\hat{v}_0\hat{U}_1^{\dagger}=\hat{X}(D-1)\otimes\hat{X}(1/3)^{\otimes 3}$. The phases $f_k(n)$ satisfy the invariance condition (2), $f_1(n)+3f_2(n)\equiv 0 \mod D$. Thus, the observable $\hat{v}_1$ has $|\psi\rangle$ as its eigenstate with eigenvalue 1. By Eq. (5) one has $\hat{X}(D-1)=\omega\hat{X}(0)$, and $\hat{Y}\equiv\hat{X}(1/3)$ is given by

$$\hat{Y}=\omega^{-1/3}\left(\sum_{n=0}^{D-2}|n\rangle\langle n+1|+\omega^{D/3}|D-1\rangle\langle 0|\right). \tag{10}$$

The observable $\hat{Y}=\hat{X}(1/3)$ is not equivalent to $\hat{X}=\hat{X}(0)$, see the explanation below Eq. (5).

The other three concurrent observables are obtained by $\hat{v}_l\equiv\hat{U}_l\hat{v}_0\hat{U}_l^{\dagger},\ l\in\{2,3,4\}$, where the unitary operators $\hat{U}_l$ are composed of the cyclic permutations of the local phase shifters: $\hat{U}_2=\hat{P}_2\otimes\hat{P}_1\otimes\hat{P}_2\otimes\hat{P}_2, \hat{U}_3=\hat{P}_2\otimes\hat{P}_2\otimes\hat{P}_1\otimes\hat{P}_2$, and $\hat{U}_4=\hat{P}_2\otimes\hat{P}_2\otimes\hat{P}_2\otimes\hat{P}_1$. The perfect correlations for $\hat{v}_i$ $(i=1,\ldots,4)$ are respectively given by

$$\begin{aligned}
&C_{\mathrm{QM}}(x_1+y_2+y_3+y_4\equiv -1),\\
&C_{\mathrm{QM}}(y_1+x_2+y_3+y_4\equiv -1),\\
&C_{\mathrm{QM}}(y_1+y_2+x_3+y_4\equiv -1),\\
&C_{\mathrm{QM}}(y_1+y_2+y_3+x_4\equiv -1),
\end{aligned} \tag{11}$$

where ω^{y_i} is an outcome of measurement $\hat{Y}$ for i-th party.

Local realistic theories assume that the outcomes of the measurements are predetermined, before the actual measurements. This means that the values of local realistic predictions for the correlations (11), for each experimental run, must satisfy:

$$\begin{aligned}
\omega^{x_1}\omega^{y_2}\omega^{y_3}\omega^{y_4}&=\omega^{-1},\\
\omega^{y_1}\omega^{x_2}\omega^{y_3}\omega^{y_4}&=\omega^{-1},\\
\omega^{y_1}\omega^{y_2}\omega^{x_3}\omega^{y_4}&=\omega^{-1},\\
\omega^{y_1}\omega^{y_2}\omega^{y_3}\omega^{x_4}&=\omega^{-1},
\end{aligned} \tag{12}$$

If local realism is also to reproduce quantum perfect correlations for the case of $\hat{v}_0=\hat{X}^{\otimes 4}$, one must have

$$E_{\mathrm{LR}}(x_1,x_2,x_3,x_4)=1, \tag{13}$$

or for each run

$$\omega^{x_1}\omega^{x_2}\omega^{x_3}\omega^{x_4}=1, \tag{14}$$

If one multiplies side-by-side all equations (12), one sees that this is possible only provided $\omega^{-3\sum_{i=1}^{4}y_i-4}=1$. As $\omega=\exp(2\pi i/D)$, if $D=3d$ where d is an integer, an elementary algebra shows that there is no integer solution of $\mathbf{y}$ of the equation $3\mathbf{y}+4\equiv 0 \mod D$. Thus local realistic correlation (14) is impossible (a GHZ-like contradiction).

It is worth noting that the approach with concurrent observables relaxes the restrictions of early studies requiring *compatible* observables. This enables one to generalize GHZ contradictions, beyond the case $N>D$ studied in Refs. [10, 12].

Note that in order to prove the 4-partite GHZ contradiction we have chosen the local dimension D and the number of the observables $\hat{Y}$'s in the considered correlation functions, which will be denoted as N_2, such that the greatest common divisor (gcd) of D and N_2, $\gcd(D,N_2)=3$, does not divide the number of parties $N=4$, or equivalently the number N_1 of the observables $\hat{X}$ (here equal to 1). This mathematical property plays a central role in the generalization of the GHZ contradiction to an arbitrary number of parties.

The unitary observables $\hat{X}$ and $\hat{Y}$ can be realized by using optical devices such as multiport beam splitters and phase shifters (see, [16]).

B. Extension to multipartite systems

We extend our approach a general case of N qu$D$its, which $N\geq 3$, which is non-divisible by any non-unit divisor of D, smaller than N.

To this end, we use a set of $(N+1)$ concurrent observables given by $\hat{v}_0=\hat{X}^{\otimes N}$ and N observables of the following forms: $\hat{v}_1=\omega\hat{X}^{\otimes N_1}\otimes\hat{Y}^{\otimes N_2}$ and $\hat{v}_k=\hat{Y}^{\otimes k-1}\otimes\omega\hat{X}^{\otimes N_1}\otimes\hat{Y}^{\otimes N_2-k+1}$ for $k=2,...,N_2+1$ and finally $\hat{v}_k=\hat{X}^{\otimes k-N_2-1}\otimes\hat{Y}^{\otimes N_2}\otimes\omega\hat{X}^{\otimes N-k+1}$ for $k=N_2+2,...,N$

The composite observable $\hat{v}_1$ is obtained by a unitary transformation, $\hat{U}_1=\hat{P}_1\otimes\mathbb{1}^{\otimes N_1-1}\otimes\hat{P}_2^{\otimes N_2}$, of the observable $\hat{v}_0$, i.e., $\hat{v}_1=\hat{U}_1\hat{v}_0\hat{U}_1^{\dagger}$, with the phase shifters $\hat{P}_1$ and $\hat{P}_2$ of phases $f_1(n)=(D-1)n$ and $f_2(n)=n/N_2$, respectively. The local observables $\hat{X}=\hat{X}(0)$ and $\hat{Y}=\hat{X}(1/N_2)$ are given by Eq. (5). Likewise, we obtain the other concurrent observables $\hat{v}_l$ $(2\leq l\leq N)$ by cyclic permutations in the unitary operator $\hat{U}_1$, as it was done in Sec. III. The phases satisfy the invariance condition in Eq. (2) as $f_1(n)+N_2f_2(n)\equiv 0 \mod D$. Thus, the N-partite generalized GHZ state $|\psi\rangle=\frac{1}{\sqrt{D}}\sum_{n=0}^{D-1}|n\rangle^{\otimes N}$ is a common eigenstate of all the $(N+1)$ concurrent observables $\hat{v}_l$ $(l = 0, \ldots, N)$, with the same eigenvalue 1. This leads to the following values of correlation functions (for later convenience we now give to the parties the indices $i = 1, ...N$, which will later on allow us to get a more concise notation in formulas). For all local observables equal to $\hat{X}$, that is global $\hat{v}_0$, one has $E_{\mathrm{QM}}(X, X, ..., X) = 1$. Thus we have a perfect correlation which can be denoted, in the way introduced earlier, as $C_{\mathrm{QM}}(\sum_{i=1}^{N} x_i \equiv 0)$. For $\hat{v}_k$, where $k = 1, 2, ...N$, one has perfect correlations of the following forms: for $k = 1$

$$C_{\mathrm{QM}}(\sum_{i=1}^{N_1} x_i + \sum_{i=N_1+1}^{N} y_i \equiv -1), \tag{15}$$

and for $k = 2, ..., N_2 + 1$

$$C_{\mathrm{QM}}(\sum_{i=1}^{k-1} y_i + \sum_{i=k}^{N_1+k-1} x_i + \sum_{i=N_1+k}^{N} y_i \equiv -1), \tag{16}$$

and finally for $k = N_2 + 2, ..., N$

$$C_{\mathrm{QM}}(\sum_{i=1}^{k-N_2-1} x_i + \sum_{i=k-N_2}^{k-1} y_i + \sum_{i=k}^{N} x_i \equiv -1). \tag{17}$$

Following the similar arguments as in the case of 4-partite GHZ contradiction, we obtain the following condition for the local realistic correlation function as for the composite observable $\hat{v}_0 = \hat{X}^{\otimes N}$, to have value equal to the quantum prediction, that is 1. It reads (modulo D)

$$N_1 \sum_{i=1}^{N} x_i \equiv -N_2 \sum_{i=1}^{N} y_i - N \equiv 0. \tag{18}$$

However, if N_2 is an integral multiple of g but N is not, i.e., $N_2 = \eta g$, and N cannot be divided by g, then there are no solutions of $\mathbf{y} = \sum_i y_i$ to the equation $N_2\mathbf{y} + N \equiv 0$ mod D. The greatest common divisor of N_2 and D is an integral multiple of g, i.e., gcd(N_2, D)=kg for some positive integer k but kg cannot divide N as N is not an integer multiple of g. Thus we have a contradiction with the quantum prediction.

In summary, in order to show a GHZ contradiction for N-partite and D-dimensional system, we choose that (a) $D = dg$, (b) $N_2 = \eta g$, where d and η are positive integers, and (c) N cannot be divided by g.

The integer g, a non-unit divisor D, plays a crucial role. For example, let us consider *four* 6-dimensional systems. The non-unit divisors g, smaller than $N = 4$, are 2 and 3. If we choose $g = 2$, then we are unable to see any 4-partite GHZ contradiction as the greatest common divisor (gcd) of N_2 and D, gcd$(N_2 = 2, D = 6) = 2$, also divides $N = 4$. If we choose $g = 3$, on the other hand, gcd$(N_2 = 3, D = 6) = 3$ but $g = 3$ does not divide $N = 4$ and the 4-partite GHZ contradiction can be proved. This is in fact a specific example of GHZ contradiction for a $(4, 2, 3d)$ problem. In other words, we are always able to prove the GHZ contradiction for $N(\geq 3)$-partite and $D(\geq 2)$-dimensional system as long as N cannot be divided by all non-unit divisors of D.

C. Comparison with previous works

For appropriate values of N and D, our approach reproduces the previous works [10, 12, 13]. The work in Ref. [10] showed a GHZ contradiction for $(D+1)$ qu$D$its. In our scheme, we are able to choose the number $N_1 = 1$ of the observable $\hat{X}$ and $N_2 = D$ of the observable $\hat{Y}$, then our GHZ contradiction for $(N, 2, D)$ system is reduced to a $(D + 1, 2, D)$ problem; $N_1 = 1$ is indivisible by $D = \text{gcd}(N_2 = D, D)$. Consider a non-unit divisor $g = 2$ of D and an arbitrary odd integer N_1. Thus we have a (odd N, 2, even D) problem, which was studied in the Refs. [12, 13]. One can also easily check that if $N_2 = D = 2$ and $N_1 = 1$, then our contradiction is reduced the original one of Greenberger, Horne, and Zeilinger [4].

IV. GENUINELY MULTIPARTITE HIGH-DIMENSION GHZ THEOREM

The GHZ theorem for 2-dimensional systems seems to be fully understood. However, for more complex systems this is not so. Cerf *et al.* suggested a criterion for a genuinely N-partite (D-dimensional) GHZ contradiction [12]. It must arise only for the given N-partite (D-dimensional) system, but not for the $n(< N)$-partite subset, or for effectively lower dimensionality of the involved observables (example: the three qubit classic GHZ theorem can put as theorem for three qutrits, and specific entangled GHZ states involving only two dimensional subspaces for each qutrit) .

The GHZ contradiction we show here is a genuinely N-partite one, as it is constructed using a set of composite observables composed of cyclic permutations. Let us explain this with the 4-partite GHZ contradiction. We use the following five concurrent composite observables:

$$\begin{gathered}
\hat{X} \otimes \hat{X} \otimes \hat{X} \otimes \hat{X} \\
\hat{X} \otimes \hat{Y} \otimes \hat{Y} \otimes \hat{Y} \\
\hat{Y} \otimes \hat{X} \otimes \hat{Y} \otimes \hat{Y} \\
\hat{Y} \otimes \hat{Y} \otimes \hat{X} \otimes \hat{Y} \\
\hat{Y} \otimes \hat{Y} \otimes \hat{Y} \otimes \hat{X}
\end{gathered} \tag{19}$$

In such a circumstance, if we eliminate one of the parties, we are not able to show a GHZ contradiction with the remaining observables. The 4-partite GHZ state is no longer their common eigenstate. Similar argument can be put forward in the case of our N-partner theorem.

The genuine D-dimensionality is reflected by the fact that the operators are undecomposable to a direct sum of any subdimensional observables [13]. In other words if two local observables $\hat{X}$ and $\hat{Y}$ can be simultaneously block-diagonalized by some similarity transformation $\hat{S}$ such that $\hat{S}\hat{X}\hat{S}^\dagger = \hat{X}_1 \oplus \cdots \oplus \hat{X}_K$ and $\hat{S}\hat{Y}\hat{S}^\dagger = \hat{Y}_1 \oplus \cdots \oplus \hat{Y}_K$, then there exist some eigenstates $|n\rangle_\alpha$ of $\hat{X}$ and $|m\rangle_\beta$ of $\hat{Y}$ such that ${}_\alpha\langle n|\hat{S}^\dagger\hat{S}|m\rangle_\beta = 0$ and one can find a sub–dimensional GHZ contradiction. However, there are no such eigenstates because for every n and m

$$|{}_\alpha\langle n|m\rangle_\beta|^2 = \frac{\sin^2(\pi\xi)}{D^2\sin^2\left[(\pi/D)\xi\right]} > 0, \tag{20}$$

where $\xi = m - n + \beta - \alpha$. As, the local observables $\hat{X} = \hat{X}(\alpha)$ and $\hat{Y} = \hat{X}(\beta)$ are such that $\beta - \alpha$ is not an integer, ξ is a non-integral rational number. Thus, our GHZ contradiction is genuinely D-dimensional.

One might try to generalize our method to more than two measurement settings. However this cannot be a genuinely multi-setting case. Consider a $(N, M, D) = (5, 3, 2)$ problem. Our approach requires six concurrent observables $\hat{v}_0 = \hat{X}\otimes\hat{X}\otimes\hat{X}\otimes\hat{X}\otimes\hat{X}$, $\hat{v}_1 = \omega\hat{X}\otimes\hat{Y}\otimes\hat{Y}\otimes\hat{Z}\otimes\hat{Z}$, and $\hat{v}_l$ formed by cyclic permutations of observables in $\hat{v}_1 (2 \le l \le 5)$. The local observables are given by the Eq. (5) with the $f_1(n) = (D-1)n$, $f_2(n) = n/8$, and $f_3(n) = 3n/8$, respectively. As the phases satisfy the invariance condition in Eq. (2) as $f_1(n) + 2f_2(n) + 2f_3(n) \equiv 0 \mod D$, the 5-partite and 2-dimensional GHZ state $|\psi\rangle = (|0,0,0,0,0\rangle + |1,1,1,1,1\rangle)/\sqrt{2}$ is a common eigenstate of these concurrent observables, with the eigenvalue equal to 1. With all that, one can show a GHZ contradiction. However, it is not a genuine 3-setting one. The contradiction also arises for observables, composed of two local measurements $\hat{X}$ and $\hat{Y}$, $\hat{u}_0 = \hat{X}\otimes\hat{X}\otimes\hat{X}\otimes\hat{X}\otimes\hat{X}$, $\hat{u}_1 = \omega\hat{X}\otimes\hat{Y}\otimes\hat{Y}\otimes\hat{Y}\otimes\hat{Y}$, and $\hat{u}_l$ formed with cyclic permutations in $\hat{u}_1 (2 \le l \le 5)$, of a similar kind as above. Thus problem whether there are genuinely multi-setting GHZ contradictions is still open.

V. SUMMARY

We gave a generalized GHZ contradiction for *multi-partite* and *high*-dimensional systems. For the purpose, we employed concurrent composite observables, which have a generalized GHZ state as a common eigenstate (even though these observables are *incompatible*). We obtained concurrent observables by using the unitary operations composed of the local phase shifters. Previous approaches had used *compatible* composite observables. Abandoning this allowed to reach a broader class of GHZ contradictions.

As a result, we showed a 4-partite GHZ contradiction for $3d$-dimensional systems, where d is an arbitrary integer. We extended the method to N-partite situation with D-dimensional systems. It turned out that non-unit divisors of D plays a crucial role to show the GHZ contradiction. We were always able to show a contradiction as long as N cannot be divided by all non-unit divisors of D, smaller than N. We also showed that our formulation of a generalized GHZ contradiction is genuinely N-partite *and* D-dimensional.

ACKNOWLEDGMENTS

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MEST) (No. 2010-0015059 and No. 2010-0018295). J.R. is supported by the Foundation for Polish Science TEAM project cofinanced by the EU European Regional Development Fund and a NCBiR-CHIST-ERA Project QUASAR. C.L. is supported by the National Research Foundation and Ministry of Education, Singapore.

[1] J. S. Bell, Physics **1**, 1 (1964).

[2] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, Phys. Rev. Lett. **23**, 880 (1969).

[3] M. Żukowski and Č. Brukner, Phys. Rev. Lett. **88**, 210401 (2002); D. Collins, N. Gisin, N. Linden, S. Massar, and S. Popescu, *ibid.* **88**, 040404 (2002); W. Laskowski, T. Paterek, M. Żukowski, and Č. Brukner, *ibid.* **93**, 200401 (2004); W. Son, J. Lee, and M. S. Kim, *ibid.* **96**, 060406 (2006); J. Lim, J. Ryu, S. Yoo, C. Lee, J. Bang, and J. Lee, New J. Phys. **12**, 103012 (2010).

[4] D. M. Greenberger, M. A. Horne, and A. Zeilinger, in *Bell's Theorem, Quantum Theory, and Conceptions of the Universe*, edited by M. Kafatos (Kluwer, Dordrecht, 1989).

[5] A. Ekert, Phys. Rev. Lett. **67**, 661 (1991).

[6] R. Horodecki, M. Horodecki, and P. Horodecki, Phys. Lett. A **222**, 21 (1996).

[7] R. Cleve and H. Buhrman, Phys. Rev. A **56**, 1201 (1997); Č. Brukner, M. Żukowski, J.-W. Pan, and A. Zeilinger, Phys. Rev. Lett. **92**, 127901 (2004).

[8] M. Żukowski, A. Zeilinger, M. A. Horne, and H. Weinfurther, Acta Phys. Pol. A **93**, 187 (1998); M. Hillery, V. Bužek, and A. Berthianume, Phys. Rev. A **59**, 1829 (1999); J. Kempe, *ibid.* **60**, 910 (1999); V. Scarani and N. Gisin, Phys. Rev. Lett. **87**, 117901 (2001); J. Barrett, L. Hardy, and A. Kent, *ibid.* **95**, 010503 (2005); A. Acín, N. Brunner, N. Gisin, S. Massar, S. Pironio, and V. Scarani, *ibid.* **98**, 230501 (2007).

[9] S. Pironio, A. Acín, S. Massar, A. B. de la Giroday, D. N. Matsukevich, P. Maunz, S. Olmschenk, D. Hayes, L. Luo, T. A. Manning, and C. Monroe, Nature **464**, 1021 (2010).

[10] M. Żukowski and D. Kaszlikowski, Phys. Rev. A **59**, 3200 (1999).

[11] D. Kaszlikowski and M. Żukowski, Phys. Rev. A **66**, 042107 (2002).

[12] N. J. Cerf, S. Massar, and S. Pironio, Phys. Rev. Lett. **89**, 080402 (2002).

[13] J. Lee, S.-W. Lee, and M. S. Kim, Phys. Rev. A **73**, 032316 (2006).

[14] J. Lee and S.-W. Lee, J. Korean Phys. Soc. **46**, 181 (2005).

[15] S.-W. Lee, Y. W. Cheong, and J. Lee, Phys. Rev. A **76**, 032108 (2007).

[16] M. Reck, A. Zeilinger, H. J. Bernstein, and P. Bertani, Phys. Rev. Lett. **73**, 58 (1994); M. Żukowski, A. Zeilinger, and M. A. Horne, Phys. Rev. A **55**, 2564 (1997).